



Royal College of
Paediatrics and Child Health

Leading the way in Children's Health

MRCPCH FACE-TO-FACE NORMAL CIRCUIT - NO PATIENTS CLINICAL EXAM

Clinical Examination Technique (Clinical Stations & Development)

Contents

FOREWORD	3
INTRODUCTION TO THE MRCPCH FACE-TO-FACE NORMAL CIRCUIT - NO PATIENTS CLINICAL EXAMINATION	5
THE CLINICAL STATION	6
General Appearance	8
Respiratory System	10
Cardiovascular System	13
Abdominal System	16
Nervous System	19
Endocrine System	24
Locomotor System	26
Skin	29
Ear Nose & Throat	30
Diagnostic Imaging	31
DEVELOPMENTAL ASSESSMENT: AGE 0 – 5 YEARS	32

Foreword

The 'Face-to-Face Normal Circuit - No Patients MRCPCH Clinical examination' has been designed to deliver a clinical examination without patients. It consists of 10 stations. All participants (examiners, candidates, role players and helpers) will be in the same location for the exam circuit. The communication, video and history stations will be unchanged. However, the 4 short clinical examination stations and the development station will NOT have patients. Instead, candidates will need to describe a virtual examination of a child similar to the way SIM training is run in Life Support Courses in the UK and overseas.

The aim of the 4 short clinical and the development station is to test the candidate's ability to describe good clinical examination technique, and to demonstrate competence in diagnostic ability and judgement. This is of central importance in clinical practice.

In these stations and within the time available, the candidate is expected to

- focus on what is most pertinent for the case
- describe the process of a focused general physical, body systems (e.g. CNS, CVS, MSK, RS) or regional (e.g. abdomen) examination or developmental assessment,
- interpret the signs and explain their significance
- solve the clinical problem and formulate the most likely differential diagnoses
- discuss the management plan that is most appropriate for the case

This is meant to replicate what the candidate would do in clinical practice during the assessment of a child. Being able to describe the relevant examination in a succinct, structured and systematic manner will be central to success. Additionally, being able to share with the examiner what you would be looking for and why will reveal your thought processes, e.g. "I will look at muscle bulk and compare both sides to see if there is wasting or hypertrophy".

In each of these five stations (ie 4 short clinical and 1 development station), when the candidate describes their process of examining a patient, the examiner will provide 'cues' at key points to help build up a clinical picture. These 'cues' are the signs the candidate would have picked up had they done that element of the examination. For instance, if the candidate said, "I will test knee reflexes by tapping on the tendon between the patella and tibial tuberosity with a hammer," the examiner would deliver a cue "knee reflex is brisk on the left".


This short booklet does not aim to present a syllabus for the clinical examination. It sets a level of technique and competence of the good candidate who meets the expected standard. The booklet is designed around the membership of the Royal College of Paediatrics and Child Health clinical examination (MRCPCH), providing guidance for examiners and candidates. It also describes what is expected in good paediatric clinical examination technique.

A valid exam reflects everyday clinical practice. The aim of the MRCPCH is to test the candidate's ability to deal with these situations. None of us would wish to learn techniques purely for the examination and then discard them in our everyday practice. This makes it essential that in the MRCPCH examination, cases selected are representative of real-life practice. The criteria used in judging candidate performance is rated against a recognised standard. The standards presented in this booklet are those of which the examiner and the candidate should be equally aware.

Anna Mathew

Introduction to the MRCPCH Face-to-Face Normal Circuit - No Patients Clinical Examination

The Clinical examination of the MRCPCH is an OSCE composed of 10 stations for each candidate.



The Clinical Station

- 4 x 9-minute stations
- Candidate will have 4 minutes of preparation time before the station starts. They will have written information that provides a brief introduction about the case and the expected task.
- Candidate will have 6 minutes to undertake task, 3 minutes discussion with examiner.
- General physical, system or regional focus in each station could be any of the following:
 - General ('head to toes'), including skin and syndromes
 - CVS
 - Neurology/Neuro-disability and including cranial nerves
 - MSK
 - Respiratory
 - Abdominal
 - Gastroenterology
 - Ophthalmology
 - Dermatology
 - Hepato-biliary and pancreas
 - Haematology
 - Nutrition, growth and endocrinology
 - Nephrology
- 1 scenario per station
- Each station scenario is reviewed by a pair of examiners and marking criteria agreed
- Examiners will explain task to candidates
- The precise mix of cases will vary across exam days
- Within the time available, the candidate who meets the expected standard will
 - focus on what is most pertinent for the case
 - describe the process of a focused examination
 - interpret the signs and explain their significance
 - solve the clinical problem and formulate the most likely differential diagnoses
 - discuss the management plan that is most appropriate for the case

- Candidates should read their Information sheet carefully and focus on the specific task for the station. The task may not necessarily require a full system examination.
- The examiners are looking for a systematic, fluid approach. The candidate who has regularly examined children during their training will be able to demonstrate this.
- Cases without abnormal physical signs may be included, with an appropriate introduction, e.g., "This young child was referred from the

community clinic. He is 16-months, and is unable to walk. What do you think?"

- Competence in examination technique, ability to elicit and interpret abnormal findings or their absence, and ability to discuss their significance will be assessed.
- When describing the process of physical examination in the child, candidates will be expected to mention the features noted in the next section (General Appearance) when examining any child.

General appearance

Irrespective of whether the station requires general physical, system or regional examination, the candidate is expected to describe general inspection and palpation that are pertinent to the patient's presentation.

If the task is general physical examination ('head to toes'), the candidate should devote more time to focused general inspection and palpation, and briefly mention other pertinent organ (e.g. palpation of liver and spleen if initial general examination reveals pallor and lymphadenopathy) or system examination (e.g. heart auscultation if short girl).

If the task pertains to system or regional examination, then the candidate should briefly describe focused general appearance and prioritise time for the system/general examination.

In order to complete all domains in the station, the candidate must be succinct, structured and systematic when describing the relevant examination. This requires the candidate to be aware of the time, to refrain from unnecessary examination and to avoid repetition.

General health

The candidate should note whether the child shows any signs of being unwell at the time of the examination. Nursing observations and growth measurements may be provided. If these are not provided but are relevant to the case, the candidate is expected to elicit this information from the examiner.

Dysmorphic features

Unusual features in the head, face, neck and limbs of a syndromic or non-syndromic pattern may be noted during system examination or the candidate may be referred back to these observations.

Growth and Nutrition

Full assessment is not usually possible in the short cases. Candidates should remember that it may be difficult to assess a child's age. The candidate should note obvious abnormalities in weight, length/height, nutritional status, adiposity and pubertal status.

Development

Neurodevelopmental abnormalities may be noted during examination of other systems.

Hands

The candidate should be able to identify finger clubbing, abnormalities of the nails, note the colour of the hands and recognise poor perfusion.

Face

Inspection should include assessment of colour, cyanosis, anaemia, jaundice and any signs of current illness or dehydration. The candidate should note craniofacial abnormalities.

General observations

Candidates may comment upon other features noted in the course of clinical examination. This may include evident clinical features, such as a boy with severe widespread eczema. Often other observations are helpful, e.g. the presence of a nasogastric tube, an intravenous infusion site, an ankle foot orthosis, an inhaler etc.

Respiratory System

Inspection

Aspects which the candidate should be able to quickly inspect include the hands, face, neck, and chest wall. (See also section on General Appearance).

Hands

The candidate should be able to identify finger clubbing and cyanosis and abnormal perfusion.

Face & neck

The candidate should be able to identify cyanosis, evidence of increased work of breathing, cough, stridor or upper airway obstructions, nasal congestion or obstruction and other signs of respiratory illness.

Chest wall

- Shape - the candidate should be able to recognise different chest wall shapes, and understand their significance (e.g. increase of AP diameter, Harrison's sulci, kyphoscoliosis, barrel chest or pectus carinatum, pectus excavatum).
- Movements - the candidate should be able to measure the rate of respiration, and assess whether accessory muscles of respiration are being used, and chest expansion is normal and symmetrical. The pattern of respiratory movement may be abnormal (e.g. asthma).
- Skin - the candidate should be able to recognise associated disease (e.g. eczema), evidence of previous surgery (e.g. thoracotomy, chest drain, Portacath) engorged superficial veins (e.g. SVC obstruction), subcutaneous emphysema.

Abdomen

The candidate should be able to comment on the movement of the abdomen with respiratory effort. Examination may include abdominal palpation (e.g. liver edge).

Palpation

Mediastinum

The candidate should be able to demonstrate appropriate techniques for assessing the position of the upper and lower mediastinum: tracheal position, chest symmetry and apex beat.

Chest wall movement

The candidate should be able to assess both the extent and the symmetry of movement of the chest wall.

Neck and axillae

The candidate should be able to palpate for cervical and axillary lymphadenopathy.

Percussion

Percussion may not be helpful in infants and toddlers. The candidate should be able to assess the percussion note over both sides of the chest, and to distinguish resonant / hyper-resonant / dull / stony dull notes.

Auscultation

Candidates should be able to elicit signs and interpret them taking account of a child's age and the presence of upper respiratory tract secretions. The candidate should be able to distinguish bronchial from vesicular breath sounds.

The candidate should be able to assess the intensity of the breath sounds, and any changes in their timing (e.g. prolonged expiratory phase in obstructive airway disease).

The candidate should be able to identify any added sounds (e.g. crackles, wheezes), and to assess their nature and timing (e.g. the fine end-inspiratory crackles).

The candidate should be able to assess vocal resonance, but only when appropriate.

Additional Points

Sputum pot

The candidate should be able to draw inferences from the contents of the sputum pot (e.g. copious mucopurulent secretions suggest bronchiectasis).

Peak flow rate

The candidate should be able to measure the peak flow rate, using the appropriate meter and know that the result relates to the child's height.

Inhaler technique

The candidate should be able to assess inhaler technique, and be familiar with standard inhaler devices and their appropriate application.

Tracheostomy

The candidate should recognise a tracheostomy or previous scar and understand why it is present.

Patterns of Abnormalities

The candidate should be able to recognise and sensibly discuss the pattern of abnormal signs which suggest the following:

- Consolidation
- Collapse or removal of a lung (or part thereof)
- Pleural effusion
- Pneumothorax
- Airflow obstruction
- Bronchiectasis

Cardiovascular System

Inspection

The candidates should be able quickly to inspect the hands, face and neck and chest wall. (See also section on General Appearance). The candidate should understand how symptoms and signs relate to age (e.g. poor feeding in the infant with heart failure, variation in pulse and BP). The effects of previous acute cardiovascular illness should be recognised (e.g. septic emboli, neurodevelopmental problems).

Hands

The candidate should be able to identify finger clubbing, cyanosis, abnormalities of peripheral perfusion, the stigmata of endocarditis and congenital hand abnormalities associated with CVS problems.

Face and Neck

The candidate should examine the lips and tongue for cyanosis. The candidate should be able to recognise abnormalities of complexion, (e.g. anaemia, polycythaemia).

Jugular Venous Pulse and Hepatomegaly

The candidate should understand examination of this aspect is very dependant upon a child's age.

The candidate should be able to differentiate arterial from jugular venous pulsation. The candidate should understand abnormalities of the jugular venous pulse in the older child, although these are rare in paediatric practice. The JVP is best seen with the child lying at 45°, with the child's head turned towards the candidate.

In the infant and young child the candidate should routinely examine the abdomen for hepatomegaly.

Chest Wall

The candidate should be able to recognise different chest wall shapes and evidence of previous surgery (e.g. median sternotomy, lateral thoracotomy, pacemaker).

Palpation

Arterial Pulse

The candidate should be aware of the components of the normal pulse waveform and their origin. The candidate should be familiar with the range of

normal heart rate at different ages. A candidate should be able to detect the following pulse abnormalities and appreciate the underlying pathophysiology:

- Sinus tachycardia
- Sinus bradycardia
- Sinus arrhythmia
- Small volume
- Radio-femoral delay or absent femoral pulses
- Collapsing
- Paradoxical pulse

The brachial pulse is suitable for assessment of the pulse at all ages. In younger children and after surgery the radial pulse may mislead. In older children, candidates may palpate the carotid pulse. Femoral pulses are mandatory.

Precordium

The candidate should attempt to locate the apex beat and be able to distinguish:

- Displacement
- Hyperdynamic apex
- Dextrocardia

The candidate should be able to test for and recognise:

- Right ventricular parasternal lift
- Palpable heart sounds
- Cardiac thrill

Blood Pressure

The candidate should be familiar with the technique of measurement of arterial pressure with a conventional sphygmomanometer, if indicated in both upper and lower limbs. The importance of cuff size should be appreciated. Candidates should relate BP to age and have a rough idea of the normal range for age. Phase IV or V recordings would be acceptable as diastolic blood pressure. The candidate should be aware of the potential significance of abnormalities in blood pressure.

Candidates should be aware of the difficulties of obtaining reliable blood pressure measurements in children. They are not expected to know how to use an automated blood pressure device.

Percussion

Percussion of the cardiac border or area of cardiac dullness adds little to the clinical assessment. Candidates are not expected to include this in routine examination.

Auscultation

Candidates should remember that the examiners will have agreed the signs that they are expected to elicit, using standards appropriate for the level of competence required.

The candidate should be able to identify normal heart sounds and the common abnormalities thereof.

- First Sounds, loud S1, split S1
- Second Sound, wide expiratory splitting, fixed splitting, single S2, loud component of S2

In addition the candidate should be able to recognise other auscultatory features such as:

- Sinus arrhythmia
- Gallop rhythm
- Heart murmur
- Ejection click
- Opening snap
- Mid systolic click
- Metallic prosthetic sounds
- Pericardial friction rub

The candidate should be familiar with the surface markings of the four valve "areas" and be able to time murmurs to diastole and systole (e.g. mid, late or pansystolic) or continuous. The loudness/intensity of the murmur should be described. The candidate may use a grading system. If appropriate, auscultation should be performed with the child in different positions.

If a murmur is heard, the candidate should also auscultate the neck and the lateral and posterior chest to assess radiation of the murmur. The candidate should be able to differentiate between innocent murmurs and murmurs related to significant valvular lesions.

Additional Points

A comprehensive examination of the cardiovascular system includes detection of sacral or lower limb oedema in the older child.

If appropriate, the candidate should seek an AV malformation and listen for a bruit.

Patterns of Abnormalities

The candidate should be able to recognise and discuss the patterns of abnormal signs. Interpretation should embrace other features (e.g. previous surgery, Down's syndrome).

- Innocent murmur
- Ventricular septal defect
- Atrial septal defect
- Pulmonary stenosis
- Pulmonary regurgitation
- Fallot's tetralogy
- Patent ductus arteriosus
- Atrio-ventricular septal defects
- Coarctation of the aorta
- Aortic stenosis
- Aortic regurgitation
- Mitral stenosis
- Mitral incompetence
- Systemic-pulmonary shunt
- Pulmonary hypertension
- Pericarditis/pericardial effusion
- Tricuspid regurgitation
- Prosthetic valve

Abdominal System

Inspection

The candidate should make a quick general assessment and briefly inspect the hands, face, upper limbs, trunk and abdomen. (See also section on General Appearance).

Face, limbs and trunk

The candidate should be able to identify anaemia, jaundice, stigmata of liver disease and should note other features associated with abnormal abdominal findings (e.g. bruising, haemodialysis, shunts).

Abdomen

- Shape - The candidate should be able to recognise different shapes of abdomen, and understand their significance, (e.g. symmetry or asymmetry, flatness or distension, prune belly, hernias).

- Skin - The candidate should be able to recognise evidence of previous surgery (e.g. herniotomy, laparotomy), monilial infection, bruising and evidence of superficial veins.
- Movement - The candidate should be able to recognise abnormal movement with respiration, visible peristalsis, or pulsations.

Palpation

The candidate should be able to differentiate normal from abnormal findings, and correctly identify the following organs:

- Liver
- Spleen
- Kidneys
- Bladder
- Female genitalia
- Male genitalia (including descent of testes)

Candidates should also mention other signs identified including abdominal masses.

Percussion

The candidate should be able to percuss in order to estimate the size of an organ (e.g. liver, spleen). The candidate should be able to examine for ascites and able to elicit shifting dullness and a fluid thrill. If necessary, the examiner can be asked to help.

Auscultation

The candidate should be able to identify:

- Bowel sounds - increased, normal, absent

Additional Points

Candidates must not perform rectal or vaginal examinations.

Candidates are not expected to examine the external genitalia or perianal region if this is likely to upset a child. In other circumstances, if this is required the candidate will be specifically asked to do this.

The candidate is expected to recognise the following:

- Nasogastric tube
- Gastrostomy
- Continuous ambulatory peritoneal dialysis or other dialysis catheter

- Ileostomy or colostomy
- Nephrostomy/vesicostomy
- Indwelling central venous access device for parenteral nutrition

Patterns of Abnormalities

The candidate is expected to recognise normal and abnormal clinical signs and to discuss the pattern of signs which suggests a diagnosis.

- Liver disease
e.g. portal hypertension, cirrhosis, storage disorder, chronic liver disease
- Splenomegaly
e.g. spherocytosis, thalassaemia, portal hypertension
- Infection
e.g. viral hepatitis, ascites, glandular fever
- Inflammatory bowel disease
e.g. Crohn's, ulcerative colitic
- Myeloproliferative disorders and haematological malignancies
e.g. leukaemia, lymphoma
- Renal disease, renal enlargement and its causes
e.g. polycystic disease, hydronephrosis and renal tumours
- Therapeutic intervention
e.g. CAPD, gastrostomy, transplant, subcutaneous infusion

Nervous System

Candidates are expected to recognise the difficulties and different approaches towards neurological examination in the co-operative 12-year-old, the playful toddler and the newborn infant. They should appreciate that often a great deal can be learnt by watching a child walk or kick a football, before reaching for the tendon hammer. An integrated response is expected combining observations of behaviour and movement with findings on traditional neurological examination. The candidate is expected to adapt all examination techniques for the child's age. (See also section on General Appearance).

Cranial Nerves

The candidate should be able to examine the cranial nerves and discuss the pattern of abnormal signs.

I. Not likely to be appropriate for examination

II. The candidate should be able to:

- Make a simple assessment of visual acuity.
- Assess vision. The candidate should be able to detect bitemporal hemianopia and homonymous hemianopia in the older child.

Eyes

Ophthalmoscopy is important but difficult in most children. A systematic approach to examination of the eye should be used. Candidates should attempt fundoscopy, using suitable techniques and recognise the limitation of their findings.

- Comment on red reflex, cornea, iris, pupil and lens.
- Candidates should recognise sunseting of the eyes.
- Assess the optic disc for optic atrophy or papilloedema.
- Assess abnormalities of the retina, including the presence of haemorrhages and abnormal retinal pigmentation.
- Recognise abnormalities of the lens (e.g. cataract) or iris (e.g. heterochromia, coloboma)

III, IV, VI the candidate should be able to:

- Make a general assessment of external ocular movements by getting a child to fix on an object and follow it in an H shaped pattern.
- Know how to test the individual oculomotor nerves.
- Examine and classify a squint and competently perform a cover test.

- Examine for nystagmus and describe and discuss any abnormality, classifying the direction of nystagmus.
- Describe and discuss the abnormal eye movements associated with poor vision.

The Pupil

The candidate should be able to examine the direct and consensual response to light and the near reaction, and be able to diagnose and discuss Horner's syndrome, third nerve palsy and congenital abnormalities (e.g. coloboma).

V. The candidate should be aware of the cutaneous distribution of the three components of the nerve, and the major muscles it innervates. They should be able to demonstrate the corneal response, the jaw jerk, the testing of masseter and temporalis, and cutaneous sensation.

VII. The candidate should be able to demonstrate and appreciate the difference between an upper motor neurone and a lower motor neurone facial palsy.

The candidate should also be aware of neurological problems associated with facial palsy (e.g. altered taste, hyperacusis, altered lachrymation).

VIII. The candidate should understand hearing testing and the most appropriate choice of method according to a child's age and neurodevelopmental status. Simple assessment by history is appropriate. They should understand when it is appropriate to perform electrophysiological tests, distraction testing, and co-operation testing and full tone audiometry.

The candidate should be able to carry out Rinné's and Weber's tests using a tuning fork.

IX. This nerve will not be tested in the examination setting.

X. The candidate should be able to recognise palatal deviation, and be aware that the palate deviates to the intact side in a unilateral palatal palsy.

XI. The candidate should test the actions of sternomastoid and trapezius.

XII. The tongue is best examined for abnormal movements as it lies at rest in the mouth. The candidate should be able to recognise and discuss:

- A unilateral palsy (ipsilateral wasting fasciculation and deviation to the paralysed side).
- Fasciculation. The examiners will have agreed that this sign is present and sufficiently obvious to be recognisable.

Higher Cortical Function

Candidates should be able to make an assessment of high cortical function:

- Speech; fluency, naming capacity, repetition and comprehension
- Orientation
- Memory

Detailed assessment of complex disorders will not form part of the examination. An ability to make a confident approach to the child with learning difficulty is important as is initial assessment of the degree of learning difficulty.

Motor Functions

The candidate should be able to assess appearance, power, muscle tone, reflexes and function. They should identify dystonia and recognise involuntary movements. Tics should be identified.

Appearance

The candidate should be able to identify muscle wasting and hypertrophy, fasciculation, movement disorders and secondary disorders (e.g. kyphoscoliosis).

Tone

Candidates should assess muscle tone. In the upper limb, the candidate should include examination of flexor tone at the elbow, and pronator tone at the wrist. In the lower limb, assessment of adductor tone at the hip, extensor tone at the knee, and plantorflexor tone at the ankle should be included. The candidate should be able to elicit spasticity, rigidity and hypotonia and interpret these findings in the light of other observation.

Power

Power should be expressed in terms of function related to the child's age. The candidate should be able to assess movements against gravity and with gravity eliminated. In the older child, a system of classification of power may be applied but this is not necessary.

Coordination

The candidate should be able to demonstrate appropriate techniques for assessing coordination in the upper and lower limbs.

Reflexes

Primitive Reflexes

The candidate should be able to elicit the grasp, rooting, Moro and asymmetric tonic neck reflexes. They should understand the normal pattern of change with age and the implications of their observations.

Tendon Jerks

The candidate should be able to elicit the biceps, supinator, triceps, knee and ankle reflexes, the abdominal responses and the plantar responses.

Sensory Function

The candidate should be able to demonstrate light-touch, proprioception, vibration sense and pin prick, providing appropriate testing materials are available.

Light-touch

The candidate should use cotton-wool and not drag the stimulus along the surface of the skin.

Proprioception

The candidate should show understanding of the sensitivity of joint position sense and be able to elicit upper limb drift with the eyes closed, or a positive Romberg's test.

Vibration test

The candidate should be given a suitable (128Hz) tuning fork.

Pin prick

The candidates are not expected to perform any procedure which may be painful or uncomfortable to the patient unless specifically invited to do so by the examiner. The candidate must be provided with appropriate, disposable sharps which do not penetrate the skin.

In testing sensation, the candidate should show understanding of:

- The need to move from areas of reduced to normal sensation when testing cutaneous sensitivity.
- The cutaneous distribution of sensory loss more commonly seen in clinical practice (e.g. cord lesion compared with peripheral nerve lesion).
- Sensory abnormalities which suggest non-organic sensory loss.

Patterns of Abnormalities

Candidates are not necessarily expected to reach a diagnosis in a child with a neurological problem suitable for short case examination. Findings should always be described and the observations related to each other. Commoner patterns of abnormality include:

- Cerebral Palsy
- Hemiplegia/quadruplegia/diplegia
- Primary muscle disorders
- Hereditary motor sensory neuropathies

Cerebellar Function

The candidate should be able to examine for nystagmus and recognise dysarthria. The candidate should be familiar with the finger-nose and heel-knee-shin tests. Dysdiadochokinesis in the upper limbs may be demonstrated in older children.

The candidates should recognise the unsteady ataxic gait and truncal ataxia. The association with abnormal head control and nystagmus (discussed previously) should be recognised. The candidate should be able to recognise more common conditions (e.g. ataxic Cerebral Palsy, ataxia-telangiectasia and Friedreich's ataxia).

Gait

Candidates should be able to observe gait abnormalities. They should ask the child to perform tasks which he/she is happy and able to perform. Ideally the child's legs should be visible and socks and shoes may need to be removed. Gait should be described and interpreted in the light of other findings (e.g. waddling gait implies pelvic girdle weakness, an important cause of which is Duchenne Dystrophy).

The candidate should be able to recognise, and sensibly discuss gaits including those associated with:

- Myopathy (waddling)
- Hemiplegia
- Spastic diplegia
- Cerebellar ataxia
- Painful limb (antalgic gait)
- Foot drop
- Trendelenburg gait

Endocrine System

Patterns of Abnormalities

Many of the techniques required to examine the endocrine system have been discussed previously. Candidates are expected to be familiar with a number of specific techniques. In many cases good observation is the key skill. (See also section on General Appearance).

Candidates should be able to examine and comment upon:

- The thyroid gland and features of thyroid disease

Candidates should be able to examine the thyroid gland and detect any focal or general abnormalities (e.g. solitary nodules or goitre). They should be aware how to inspect the gland, to observe during swallowing and to palpate from behind the child if possible. Attention should be paid to related structures and tracheal displacement. Candidates should be able to assess overall thyroid status and note the features of dysthyroid eye disease.

Growth

Candidates are expected to be able to assess a child with growth problems. They should assess growth on an appropriate chart which the candidate completes or on which the child's growth measurements have been entered previously. They should demonstrate familiarity with inherited growth patterns, height, length, weight and head circumference and understand growth velocity charts. Candidates are also expected to make a brief assessment of a child's height and weight for age on simple inspection. They should note variations in normal growth and abnormal patterns of growth including those without a primary endocrine cause, hormone deficient states and growth abnormalities associated with recognised syndromes (e.g. Marfan's syndrome, Cornelia de Lange, Turner's syndrome, achondroplasia, Russell Silver syndrome).

Pubertal status/genitalia

Candidates are advised not to examine the genitalia if this would embarrass or upset the child. If appropriate, examination of the female genitalia is by simple inspection. The male genitalia should be inspected, common abnormalities of the penis recognised and descent of the testes determined.

The candidate should be able to assess pubertal status on simple examination using the Tanner stages. Approximate assessment of testicular volume may be expected but the candidates are not expected to use an orchidometer.

The child with diabetes

The candidate should understand that in most children with diabetes there will not be abnormal findings. They should know how to examine injection sites, note lipoatrophy, lipohypertrophy and be aware of rare complications of diabetes.

Adreno-cortico insufficiency

Most children will not have signs. Candidates should recognise the features of hypoadrenalism. They should know to look for pigmentary changes and hypotension in Addison's disease. They should be familiar with the features of congenital adrenal hyperplasia.

Cushingoid features

Candidates should be able to recognise the features of primary or secondary excess of corticosteroids. They should comment upon abnormal distribution of fat, striae, hypertension, abnormal facies and adrenal virilism.

Obesity

Candidates should be able to assess simple obesity and specific endocrine disorders (e.g. cushingoid features) and understand the different growth features.

Disorders of lipid metabolism

The candidate will be expected to be aware of the clinical manifestations of abnormal lipid disorders (e.g. premature arcus, xanthelasma and xanthomata).

Locomotor System

A clear introduction to the child will be given by the examiners. A structured approach is required to include an assessment of gait, arms, legs and spine (see also section on General Appearance).

General Locomotor Examination

Gait

The candidate should be able to differentiate by inspection normality from abnormality, and to characterise abnormal gait patterns (as above).

Joint appearance and movement

The candidate should be able to differentiate by inspection normality from abnormality and to specify abnormality with respect to swelling, deformity, restricted movement. They should note number and distribution of joints involved in arthritis and understand the significance of these findings. They are expected to be able to examine for associated abnormalities (e.g. psoriasis).

Muscle wasting

The candidate should be able to differentiate by inspection normal and wasted muscles, to differentiate global from localised wasting, and to specify involved muscles or groups.

Function

The candidate should be able to assess function with relation to activities of daily living e.g. mobility and dressing.

Examination of Individual Regions

The candidate should look, feel, and assess active and passive movement of joints in the examination of hands, elbows, shoulders, spine (cervical, thoracic, lumbar), hips, knees, feet and ankles. The examination should be based on the “look, feel, move”, approach to each joint and function may be assessed (e.g. gait for hip, knee and foot/ankle). The candidate should be able to:

- Detect abnormalities at these regions.
- Differentiate joint disease from periarticular lesions.
- Define and describe joint abnormality in terms of joint inflammation and/or damage.

The candidate should be able to detect the following signs at non-axial joints: increased warmth, swelling (fluid, soft tissue, bony) fluctuance, joint-line tenderness, coarse crepitus, restriction of movement, stress pain, associated muscle wasting and weakness. The candidate should be able to recognise the

associated systemic and multisystem feature of arthritis and connective tissue disease and the need to assess other systems as appropriate.

Patterns of Abnormalities

The candidate should be able to recognise the clinical presentation, and compose an appropriate differential diagnosis for the following:

- Acute monoarthritis
- Chronic monoarthritis, Oligoarthritis and Polyarthritis
- Scoliosis
- Dislocated hip and developmental dysplasia at the hip
- Contracture syndromes including arthrogyrosis
- Congenital deformities
- Functional gait abnormalities
- The limping child

The role of a musculoskeletal screening examination (pGALS)

- The pGALS screening examination (paediatric Gait, Arms, Legs and Spine) is simple and quick and helps to localise the site of joint problems.
- pGALS is very useful to identify the pattern of joint involvement especially where symptoms are illocalised
- The pGALS screen findings help to focus a more detailed regional examination (as given above) of the affected joint(s)
- An example of use of the pGALS screen follows;

The examiner at the Musculoskeletal / Other station tells you;

“This mother has noticed a problem with her child’s walking - please comment on what you see and examine the child appropriately.”

The child is sitting in a chair and you notice a swollen knee and thigh muscle wasting and tell the examiner this.

Firstly, check the child is not in pain and before requesting the child to walk and observe for limp. Then you request the child to lie supine on the couch and focus your examination on the child’s legs with inspection, palpation, movement of the knee and assess for leg length discrepancy, Baker’s cyst and quadriceps wasting.

You present your findings at the knee and comment on the gait.

The examiner asks you to comment on the differential diagnosis - and you list the causes of a swollen joint, comment on red flags to suggest sepsis and any features to suggest chronicity.

The examiner asks you what further aspects of the clinical assessment are needed.

You would be expected to comment on what the history (e.g. pointers to infection, reactive arthritis, inflammatory arthritis, and trauma) and be expected also to comment on the importance of a pGALS screen to assess for joint involvement elsewhere which may influence the differential diagnosis.

The examiner may ask you to perform the pGALS screen or a part of it and may ask what in particular you would be looking for – e.g.

- *cervical spine, symmetrical hip and ankle joint involvement (polyarticular Juvenile Idiopathic Arthritis)*
- *asymmetrical finger and toe joint involvement and psoriasis changes to nails or skin patch (psoriatic arthritis)*
- *thin, small child with arthritis at the hip and knee (inflammatory bowel disease and associated chronic arthritis)*
- *scoliosis (in the child with a leg length discrepancy)*

Skin

The candidate should be able to:

Identify and describe individual skin lesions, e.g.:

- Macules
- Papules
- Vesicles
- Purpura
- Telangiectasia
- Haemangioma
- Other congenital naevi

Recognise common or important skin disorders, e.g.:

- Eczema
- Psoriasis
- Urticaria
- Cutaneous candidiasis
- Herpes zoster
- Dermatitis herpetiformis

Identify and sensibly discuss the significance of certain skin manifestations, e.g.:

- Erythema nodosum
- Butterfly rash
- Lymphoedema
- Hyperpigmentation
- Hypopigmentation

Identify and sensibly discuss abnormalities of nails or hair:

- Onycholysis
- Nail pitting
- Alopecia

Ear Nose & Throat

The candidate is expected to make a competent examination of the ear, nose and throat and associated lymph nodes. (See also section on General Appearance).

Ears

The candidate should direct the child's parents to assist in holding the child appropriately while the ears are examined. The auriscope should be held in such a way that movement of the child does not allow injury to the external meatus. Candidates should be able to recognise conditions including otitis externa, ear wax, otitis media, chronic serous otitis media (glue ear), perforated ear drum, grommets.

Nose

The examination of the nose is simply by inspection and observation. The candidate should know how to examine the anterior nares with an auriscope.

Throat

Assisted by the parents in the younger child, the candidate should know how to examine a child's posterior pharynx and tonsillar region. In the co-operative child this should be achieved without causing the child to gag. The candidates should recognise conditions such as acute inflammation, tonsillar hypertrophy and congenital abnormalities.

Diagnostic Imaging

Candidates are expected to examine and interpret plain radiographs and other frequently used diagnostic images. This can be used in the context of a child who is being examined or in isolation. Selection of diagnostic images should emphasise the techniques which are used in acute medicine and surgery. Plain radiography is ideal. The candidates are expected to understand other commonly used diagnostic techniques.

The candidate should be able to recognise the type of diagnostic image. They should demonstrate a systematic approach to the interpretation and be aware of the limitations of the different techniques. They should be able to recognise and interpret diagnostic images at the level which one would expect of a new specialist registrar.

SYSTEM	SUITABLE	UNSUITABLE
Respiratory	chest x-ray CT	bronchogram VQ scan
Cardiovascular	chest x-ray	ultrasound catheter studies arteriography
Abdominal	abdominal x-ray	ultrasound
Renal	abdominal x-ray DMSA/Mag 3 MCUG CT/MR	ultrasound intra-operative techniques IVU
Liver	Hida or equivalent abdominal x-ray	ultrasound ERCP
Gastrointestinal	abdominal x-ray contrast swallow/meal contrast enema isotope Meckel's scan	endoscope radiology isotope studies
Neurology	CT/MR skull x-ray neonatal brain ultrasound	contrast myelography arteriography isotope studies
Bones and joints	plain radiology isotope bone scan	arthrography ultrasound bone age

Developmental Assessment: Age 0 – 5 years

- 1 x 23-minute station
- Candidate will have 10 minutes to gather history from the role player, with 2 further minutes to briefly summarise the important points to the examiner
- Candidate will have 8 minutes to describe their approach to specific aspects of developmental assessment, briefly summarise the important points, give their conclusion about the developmental age and discuss the likely explanations for the developmental problem.
- Candidate will have 3 minutes to discuss management planning with the examiner.
- The focus for developmental assessment in each station could be any of the following:
 - **Gross motor** - posture and movement
 - **Fine motor** - vision and manipulation
 - **Speech and language** - hearing and speech
 - **Social and Communication skills**
- 1 scenario per station
- Each scenario is reviewed by a pair of examiners and marking criteria are agreed.
- Examiners will explain task to candidates
- The precise mix of cases will vary across exam days
- Within the time available, the candidate who meets the expected standard will
 - focus on what is most pertinent for the case and their development problems
 - describe the process of a focused development assessment
 - interpret the findings, explain their significance and give their conclusion about the developmental age
 - discuss the most likely explanations or differential diagnoses
 - discuss the management plan that is most appropriate for the case

Full developmental assessment is not possible within the time available in the Development Station and candidates will be presented with 1 or at most 2 areas of developmental assessment to undertake.

Candidates have 23 minutes in this station, which is divided into 2 parts. The first part will be to undertake a focused developmental history and the 2nd part will be to describe the process of the developmental assessment posed by the examiner.

Candidates should use information from any universal cues provided in the station to help build the clinical picture. Examiners will then provide further dependant cues according to the assessment process the candidate describes.

Candidates will be expected to offer a view on the child's neurodevelopmental status. Candidates should be expected to assess development in children between 0-5 years of age. (See also section on General Appearance).

Candidates should be confident and proficient in assessing development in these four categories:

- **Gross motor** - posture and movement
- **Fine motor** - vision and manipulation
- **Speech and language** - hearing and speech
- **Social and Communication skills**

They should understand the confidence that they should place on their findings. There is no prescribed set of developmental tests. Use of any set of appropriate milestones is acceptable.

Within each developmental category it should be possible to define developmental age within 2-3 months before 2 years of age, and within 4-6 months between 2 and 5 years of age.

The candidates should present their findings based on the specific task presented to them. The estimated developmental age should be supported by evidence of the age-appropriate skills which the child can perform and the observation that a child is not able to perform a skill which might be expected of a slightly older child.

Interpretation of the developmental assessment should be made with regard to the range of normal findings and in the context of a child's illness or other associated condition. If neurodevelopmental delay is suggested, the candidate should be able to comment on the confidence and significance of this finding.

In the Development station the candidate should be able to discuss appropriate management steps for the child with developmental problems.

A few things to note:

The developmental toys will be available from the very beginning of the Development station and will remain available throughout the station.

*While taking the history from the role player, Candidates will be expected to **NOT** ask detailed questions about the specific development which is the focus of the station. For example,*

- *if the focus is Fine motor assessment, the candidate would be expected to ask about vision **BUT** should **NOT** ask the role player specific questions about fine motor skills (e.g. "does she pick up tiny objects and when did she first start to do this?")*

- *if the focus is Speech & Language assessment or Social & Communication skills, the candidate would be expected to ask about hearing BUT should **NOT** ask detailed questions about verbalisation (e.g. "does he babble and when did he start babbling?") or feeding (e.g. "can he feed himself with a spoon?"*
- *if the focus is Gross motor, candidates would be expected to avoid asking the role player questions such as "when did your child start sitting, standing, crawling walking, etc." Such detailed inquiries should be reserved for the second part of the station, when candidates will be able to elicit relevant information from the examiner while they describe their approach to developmental assessment.*

Examples of Scenarios generated for this station

This station examines the candidate's ability to assess specifically requested areas in a child with a developmental problem.

This may be a child with a neurological problem or syndrome who is developmentally delayed, or it may be a child who has an abnormal pattern of development e.g. autistic spectrum disorder.

Example 1

4-year-old with right hemiplegia. Please gather the relevant Developmental history from the Role player and with the examiner describe the assessment of his **fine motor skills**.

Examples of tools that might be considered:

- 12 x 1-inch blocks
- **Scissors**
- **Colouring pencils and paper**
- **Small threading beads**
- **Picture book**

What might be expected:

To the role player: Information gathering/Developmental History.

To the examiner: Description of the Assessment of building blocks skills 12 block tower or patterns of three steps using 6 blocks or more (9-10 blocks and can copy a 3-block pattern at age 3)

Assessment of whether the child can cut paper (age 3)

Assessment of whether the child can draw a man with head, body, legs and arms

Assessment of whether the child can copy an X, V, H, T and O?

Assessment of whether the child can write his or her name

Assessment of whether child can lace small beads (large at 3)

Assessment of whether child can turn the pages of a book

Assessment of whether the child can perform well using both hands? – would be looking for the functional use of right hand – is it a prop? Can the child open a yogurt or a packet of crisps?
Vision – does he have a field defect obviously impairing fine motor skills?

Example 2

3-year-old girl with Downs Syndrome. Please gather the relevant Developmental history from the Role player and with the examiner Describe a Speech and Language assessment.

Tools may include small everyday objects and pictures

What might be expected:

To the role player: Information gathering/Developmental History.

To the examiner: Enquire, first cooing, babbling, words – when was first word with meaning, is she putting 2 words together – explore to ensure whether she is putting 2 words together, concerns about hearing, ENT interventions

Assessment of concentration and attention

Assessment and understanding

Following a one or two stage command

Does she know body parts?

Assessment of object recognition and selection

Assessment of picture recognition and selection

Imitation of sounds and words

Words together – noun phrases and verb phrases

Example 3

4-year-old child whose sibling has ASD. Please gather the relevant Developmental history from the Role player and with the examiner Describe AN assessment to determine whether you think it is likely that this child has ASD

Tools may include a range of toys, ball and pretend play toys

What might be expected:

To the role player: Information gathering/Developmental History.

To the examiner: Assessment of Social communication

Communication: speech history, would listen for anything unusual about way he speaks, learned phrases, socially inappropriate questions, pronoun reversal, pretend play and imitative play

Social interaction: Assessment of interpersonal communication – does he point?

Does he take mother to what he wants? Does he share toys? How is his eye contact, does he prefer to play on his own? Does he get emotional when his mother does? Use of gestures? Friends?

Stereotyped behaviour: Assessment of ritualistic behaviour –does he like spinning, is he obsessive, rigid, routines? Assessment of other traits – does he dislike crowded spaces; does he dislike loud noise? Does he dislike having his hair cut or washed?

Example 4

18-month-old with development delay. Please gather the relevant Developmental history from the Role player and with the examiner Describe your assessment of **fine motor skills**.

Tools may include: small objects and toys – bricks, ball, doll, rattle, small picture book, in/out container, crayon/paper

What might be expected:

To the role player: Information gathering/Developmental History.

To the examiner:

Assessment of grasp – scissors or pincer grasp

Assessment of pointing – with index finger at objects of interest

Assessment of release of a small object into someone's hand

Assessment of crayon grasp and scribble - Assessment of turning of pages of a book

Build two brick towers (18-month-old should be able to build 3 or more)

Example 5

4-year-old with spastic diplegia GMFCS. Please gather the relevant Developmental history from the Role player and with the examiner Describe your assessment of **gross motor skills**

Tools may include: space to walk, stairs

What might be expected:

To the role player: Information gathering/Developmental History.

To the examiner:

History: pregnancy and birth, motor development (sitting, cruising, standing on tip toes, cruising). Enquire about upper limb function, use of a wheelchair if tired, how far can child walk?

Can the child walk, with and without shoes? any walking aids and of shoes.

Enquiry about skills on other surfaces, of ability on stairs (if unable to provide) and in playground, whether he/she can ride a tricycle.

Example 6

3-year-old with some loss of visual acuity (could be any age). Please gather the relevant Developmental history from the Role player and with the examiner Describe your assessment of this child's functional vision.

Tools may include: Black and white picture boards, keeler cards if possible, books/boards with clear pictures that can be held at a distance to functionally assess what can be seen from a distance. Small coloured beads or bricks to name colours. Hundreds and thousands and smarties to assess visual acuity of small objects. Fonts of different sizes.

What might be expected:

To the role player: Information gathering/Developmental History.

To the examiner:

Fixing and following? Response to light – blinking in sun or to light being put on. Object recognition, face recognition (no sound), in isolation and picking a face out of a group. Bumping into things, tripping up. Sitting close to the TV, use of a touch screen

Reference:

A good reference is "Child Development. An illustrated guide by Carolyn Meggitt and Gerald Sunderland (ISBN 0-435-42056-9) published by Heinemann Educational Publishers, which outlines normal development at each key stage. Details on what is expected of candidates when performing clinical examination of children are available on the RCPCH website.

Tips

Follow a clear and systematic pattern - for example if candidates are asked to Describe the fine motor skills assessment of a three-year-old candidates may decide to start with building blocks, building towers and bridges. A candidate might then move on to using crayon or pencils and paper - copy a straight line, then a circle, then a T, H, V or an X. Can the child draw a man? Candidates might choose threading beads and see how the child copes and next see how they use scissors.

For gross motor skills sitting, standing, walking, running, hopping, jumping, throwing and catching a ball may be enquired about.

What might examiners be looking for in a developmental history?

Examples of things examiners might expect include:

- Exploration of likely aetiology: perinatal history, prematurity, birth history, chromosomal disorders.
- Past history: CNS infection, trauma, malignancy.
- FH of developmental delay/learning difficulties.
- Delay in other domains or isolated issue.
- Associated comorbidity: ADHD, Epilepsy, ASD.
- Impact of child's difficulties on family: social support, income support, additional help.
- Education plan: EHCP, mainstream/special needs school.
- Team involved in child's care: MDT etc.

Candidates can make notes during the station, but they must destroy these when they finish.

It is not the examiners responsibility to remind candidates on how much time is left throughout the station. Candidates will receive a warning with 4 minutes left to go in the history plus the standard warning at 3 minutes remaining with examiner (after 20 minutes).